

# CHILD IMMUNIZATION PROGRAMME



# برنامج تحصين الطفل

العمر Age	اللقاحات Vaccines
عند الولادة Birth	BCG, Hepatitis B (1)
شهران 2 months	DTaP/Hib/Polio (1) & Hepatitis B (2) Pneumococcal (1) Check BCG scar Rotavirus *vaccine*
٤ أشهر 4 months	DTaP/Hib/Polio (2) & Hepatitis B (3) Pneumococcal (2) Rotavirus *vaccine*
٦ أشهر 6 months	DTaP/Hib/Polio (3) & Hepatitis B (4) Pneumococcal (3) Rotavirus Vaccine* Flu Vaccine**
٩ أشهر و أكثر 9 months onwards	Meningococcal ***
أشهر 12 months	MMR (1) & Chickenpox (1)
١٢ - ١٣ شهراً و أكثر 12 - 13 months onwards	Pneumococcal (4) Hepatitis A (1)****
١٨ شهراً 18 months	DTaP/Hib/Polio (4) & MMR (2)
من عمر السنتين From 2 years	(يمكن أخذه كل ٣ سنوات) Typhoid vaccine (may repeat every 3 years)
٥ - ٦ سنوات 5-6 years	DTaP/Polio (5) & Chickenpox (2)
من ٩ سنوات فما فوق From 9 years onwards	Human Papilloma Virus Vaccines (Cervical / Penile Cancer)  9-14 years : 2 doses (0, 6-12 months) 15-45 years : 3 doses (0, 2, 6 months)
١٠ - ١٢ سنوات 10-12 years	TdaP or Td

\* Rotavirus Vaccine : Given in 2 doses if Rotarix or 3 doses if Rota Teq

\*\* Flu Vaccine can be given from age of 6 months: initially 2 doses 4-6 weeks apart then annually before the winter season, starting from September.

\*\*\* Meningococcal Vaccine: from 9-23 months given in 2 doses 3 months apart or as a single dose from 2 years onward.

\*\*\*\* Hepatitis A: Given from age of 1 year in 2 doses 6-12 months apart.